

# Network Algorithms and Dynamics

Homework 1

Due: 02/08/2017

1. Consider the Chernoff bound for the summation of  $n$  iid random variables  $X_1, \dots, X_n$ , i.e.,

$$P\left(\sum_{i=1}^n X_i \geq na\right) \leq e^{-nh(a)}.$$

Prove that  $h(a) > 0$  if  $a > E[X_1]$  and  $E[e^{\theta X_1}] < \infty$  over  $\theta \in [0, b]$  for some  $b > 0$ .

2. Find a Chernoff bound for the summation of  $n$  iid random variables  $X_1, \dots, X_n$  in the form

$$P\left(\sum_{i=1}^n X_i \leq na\right) \leq e^{-n\tilde{h}(a)},$$

and identify  $\tilde{h}(a)$ .

3. Let  $X$  be the number of heads in a sequence of  $n$  independent fair coin flips, and  $\mu = E[X] = \frac{n}{2}$ . Apply the Chernoff bound to show that for any  $\delta > 0$ ,

$$P(|X - \mu| > \delta\mu) \leq 2 \exp\left(-\frac{\mu\delta^2}{3}\right).$$

Then use this bound to show that the deviation from the mean  $\mu$  is of the order  $O(\sqrt{n \log n})$ .

4. Consider the exponentially tilted distribution  $\{\hat{p}_k(s)\}$ ,  $s \in \mathbb{R}$ ,

$$\hat{p}_k(s) = p_k \frac{s^k}{\phi(s)}, k = 0, 1, \dots,$$

where  $\phi(s)$  is the generating function of  $\{p_k\}$ . What is the mean of  $\{\hat{p}_k(s)\}$ ?

5. Geometric branching process: Consider a Galton-Watson branching process with geometric offspring distribution with parameter  $p$ , i.e.  $P(\xi = k) = p^k(1 - p)$ .

(i) Compute the probability of extinction occurring in generation  $n$  (using generating functions).

(ii) Give a general expression for the probability of extinction.

(iii) Derive an expression for the generating function of

$$W = \lim_{n \rightarrow \infty} \frac{X_n}{(E(\xi))^n},$$

where  $X_n$  is the number of individuals in generation  $n$ .